

## Orthopaedic Surgeon, Foot & Ankle & Trauma Surgery

Clinical Condition	Ankle/Hindfoot Arthritis
Eligibility	Pain around the ankle (Ankle, Subtalar, Talonavicular or Calcaneocuboid joint) Associated swelling and deformity Walking is difficulty especially on uneven ground Usually Unilateral Usually a history of trauma Associated inflammatory arthropathy
Clinical photo	
Differential Diagnosis	<ol> <li>Tibialis posterior tendon insufficiency</li> <li>Fracture around the ankle</li> </ol>
Information required with referral	History: Detail history of the pain ( Location, severity, type, aggravating and relieving factors ) Range or movement of the joint Associated medical conditions Any treatment used ( orthotics , brace, analgesia) Exact location of the swelling Associated deformity
Investigation required for the diagnosis	Weight bearing Ankle x rays (AP, Lat and Mortise view) Weight bearing Foot x rays (AP, Lat and oblique view)
Pre-Referral management strategies (information required with referral)	Analgesic +/- anti inflammatory medication Activity modification Walking stick/ frame Appropriate footwear Orthotics/ brace
Referral criteria	Continuous pain despite the non-operative management.
Fact sheets	http://www.footeducation.com/subtalar-arthritis http://www.footeducation.com/rheumatoid-arthritis-talonavicular- joint http://www.footeducation.com/ankle-arthritis